



Ceftazidime-Avibactam as Salvage Therapy for Infections Caused by Carbapenem-Resistant Organisms

Elizabeth Temkin,^a Julian Torre-Cisneros,^{ij} Bojana Beovic,^b Natividad Benito,^{c,d} Maddalena Giannella,^e Raúl Gilarranz,^f Cameron Jeremiah,^g Belén Loeches,^h Isabel Machuca,^{ij} María José Jiménez-Martín,^k José Antonio Martínez,^l Marta Mora-Rillo,^h Enrique Navas,^m Michael Osthoff,ⁿ Juan Carlos Pozo,^o Juan Carlos Ramos Ramos,^h Marina Rodríguez,^o Miguel Sánchez-García,^k Pierluigi Viale,^p Michel Wolff,^{q,r} Yehuda Carmeli^{a,s}

Department of Epidemiology and Preventive Medicine, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel^a; Department of Infectious Disease, University Medical Center Ljubljana, Ljubljana, Slovenia^b; Infectious Diseases Unit, Department of Medicine, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain^c; Institut d'Investigació Biomèdica Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain^d; Infectious Diseases Unit, Department of Medical and Surgical Sciences, S. Orsola-Malpighi Hospital, University of Bologna, Bologna, Italy^e; Department of Clinical Microbiology, Hospital Universitario de Gran Canaria Doctor Negrín, Las Palmas de Gran Canaria, Spain^f; Department of Infectious Diseases, St Vincent's Hospital, Melbourne, Australia^g; Infectious Diseases Unit, Hospital Universitario La Paz-IdiPAZ, Madrid, Spain^h; Department of Infectious Diseases, Hospital Universitario Reina Sofía, Córdoba, Spainⁱ; Instituto Maimónides de Investigación Biomédica, Universidad de Córdoba, Córdoba, Spain^j; Critical Care Department, Hospital Clínico San Carlos, Madrid, Spain^k; Department of Infectious Diseases, Hospital Clínic, IDIBAPS, Barcelona University, Barcelona, Spain^l; Infectious Diseases Department, Hospital Ramón y Cajal, Madrid, Spain^m; Division of Infectious Diseases and Hospital Epidemiology, University Hospital Basel, Basel, Switzerlandⁿ; Department of Critical Care Medicine, Hospital Universitario Reina Sofía, Córdoba, Spain^o; Department of Medical Surgical Sciences, Alma Mater Studiorum University of Bologna, Bologna, Italy^p; Centre Hospitalier Universitaire Bichat-Claude Bernard, AP-HP, Paris, France^q; Université Paris Diderot, Paris, France^r; Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel^s

ABSTRACT Ceftazidime-avibactam (CAZ-AVI) is a recently approved β -lactam- β -lactamase inhibitor combination with the potential to treat serious infections caused by carbapenem-resistant organisms. Few patients with such infections were included in the CAZ-AVI clinical trials, and clinical experience is lacking. We present a case series of patients with infections caused by carbapenem-resistant *Enterobacteriaceae* (CRE) or *Pseudomonas aeruginosa* (CRPa) who were treated with CAZ-AVI salvage therapy on a compassionate-use basis. Physicians who had prescribed CAZ-AVI completed a case report form. We used descriptive statistics to summarize patient characteristics and treatment outcomes. We used the Wilcoxon rank sum test and Fisher's exact test to compare patients by treatment outcome. The sample included 36 patients infected with CRE and two with CRPa. The most common infections were intra-abdominal. Physicians categorized 60.5% of patients as having life-threatening infections. All but two patients received other antibiotics before CAZ-AVI, for a median of 13 days. The median duration of CAZ-AVI treatment was 16 days. Twenty-five patients (65.8%) concurrently received other antibiotics to which their pathogen was nonresistant *in vitro*. Twenty-eight patients (73.7%, 95% confidence interval [CI], 56.9 to 86.6%) experienced clinical and/or microbiological cure. Five patients (20.8%) with documented microbiological cure died, whereas 10 patients (71.4%) with no documented microbiological cure died ($P = 0.01$). In three-quarters of cases, CAZ-AVI (alone or combined with other antibiotics) cured infections caused by carbapenem-resistant organisms, 95% of which had failed previous therapy. Microbiological cure was associated with improved survival. CAZ-AVI shows promising clinical results for infections for which treatment options are limited.

KEYWORDS carbapenem resistance, case series, ceftazidime-avibactam

Received 13 September 2016 **Returned for modification** 29 September 2016 **Accepted** 8 November 2016

Accepted manuscript posted online 28 November 2016

Citation Temkin E, Torre-Cisneros J, Beovic B, Benito N, Giannella M, Gilarranz R, Jeremiah C, Loeches B, Machuca I, Jiménez-Martín MJ, Martínez JA, Mora-Rillo M, Navas E, Osthoff M, Pozo JC, Ramos Ramos JC, Rodríguez M, Sánchez-García M, Viale P, Wolff M, Carmeli Y. 2017. Ceftazidime-avibactam as salvage therapy for infections caused by carbapenem-resistant organisms. *Antimicrob Agents Chemother* 61:e01964-16. <https://doi.org/10.1128/AAC.01964-16>.

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Yehuda Carmeli, yehudac@tlvmc.gov.il.

Ceftazidime-avibactam (CAZ-AVI), a β -lactam- β -lactamase inhibitor combination, was approved by the U.S. Food and Drug Administration (FDA) in February 2015 (1). It is indicated for the treatment of complicated urinary tract infection (cUTI) and complicated intra-abdominal infection (cIAI) (in combination with metronidazole) in adults with limited or no other therapeutic options. A promising characteristic of CAZ-AVI is its potential to treat infections caused by carbapenem-resistant *Enterobacteriaceae* (CRE) or carbapenem-resistant *Pseudomonas aeruginosa* (CRPa). Avibactam recovers the activity of ceftazidime by inhibiting Ambler class A, class C, and some class D beta-lactamases, including the KPC and OXA-48 carbapenemases; CAZ-AVI is not active against metallo- β -lactamases (MBLs), such as NDM, IMP, and VIM (2, 3). In an *in vitro* study that included 276 meropenem-nonsusceptible *Klebsiella* species isolates, 98.9% were susceptible to CAZ-AVI (4). In a second study, 67.4% of 396 meropenem-nonsusceptible *P. aeruginosa* isolates were susceptible to CAZ-AVI (5). A third study reported 100% susceptibility to CAZ-AVI among 133 non-carbapenemase-producing *Enterobacteriaceae* isolates in which the mechanism of carbapenem resistance was extended-spectrum beta-lactamase (ESBL) and/or AmpC production together with porin deficiency (6).

Clinical data on the efficacy of CAZ-AVI against carbapenem-resistant organisms in humans are scarce. In phase II trials (7, 8) and recently completed phase III trials (9, 10) of CAZ-AVI to treat cUTI and cIAI, the comparator drug was a carbapenem; therefore, patients with infections caused by carbapenem-resistant organisms were excluded. In a pathogen-directed open-label phase III trial (the REPRISE trial) comparing CAZ-AVI to the best available therapy for treatment of cUTI and cIAI caused by ceftazidime-resistant Gram-negative organisms (11), few patients with carbapenem-resistant infections met the trial's inclusion criteria. Among the 292 *Enterobacteriaceae* isolates recovered from 288 patients in the REPRISE trial, only nine harbored non-MBL carbapenemases: six KPC-producing *Klebsiella pneumoniae* isolates and three OXA-48-producing *K. pneumoniae* isolates (12). In an effort to amass data about the effectiveness of CAZ-AVI against carbapenem-resistant organisms, we present a case series of patients with infections caused by CRE or CRPa who were treated with CAZ-AVI salvage therapy on a compassionate-use basis. Our primary aim was to evaluate three outcomes: clinical cure at the end of treatment, microbiological cure at the end of treatment, and all-cause in-hospital mortality. Our secondary aim was to identify predictors of cure and of survival.

RESULTS

Sample. Of the 25 physicians contacted, 17 responded and 15 contributed data for at least one patient with an infection caused by CRE or CRPa that was sensitive to CAZ-AVI, for a total of 38 patients. Included patients had been treated in Europe and Australia in the years 2013 to 2016. Fifteen patients came from a single hospital in Spain that had an outbreak of KPC-producing *K. pneumoniae*; the characteristics of this strain have been previously described (13). Thirty-six patients (94.7%) received CAZ-AVI as salvage therapy after treatment with other antibiotics had failed; in the other two patients, CAZ-AVI was the first antibiotic chosen because no other appropriate treatment was available. Three patients were treated at institutions where CAZ-AVI clinical trials were conducted, but they were treated on a compassionate-use basis because they met the trial's exclusion criteria.

Organisms. Thirty-four patients were infected with *Klebsiella pneumoniae*, one with *Klebsiella oxytoca*, one with *Escherichia coli*, and two with *P. aeruginosa*. Antimicrobial susceptibilities are presented in Table 1. All isolates but one were classified by the local laboratories as resistant to imipenem. In the patient with an imipenem-susceptible carbapenemase-producing organism (OXA-48-producing *E. coli*), treatment with imipenem had resulted in microbiological failure. All isolates were resistant to ceftazidime alone, with MICs ranging from 8 to ≥ 64 $\mu\text{g/ml}$. Only 14 of 34 isolates tested (41.2%) were susceptible to colistin. Table 2 summarizes the MICs of carbapenems in the 33 isolates for which carbapenem susceptibility was reported quantitatively. One patient

TABLE 1 Antimicrobial susceptibility of isolates from patients with carbapenem-resistant infections treated with compassionate-use CAZ-AVI

Antibiotic	No. of isolates tested ^a	% Susceptible
Imipenem	36	2.8 ^b
Meropenem	33	0.0
Ceftazidime	38	0.0
Colistin	34	41.2
Gentamicin	37	51.4
Amikacin	38	31.6
Tigecycline	32	62.5
Fosfomycin	29	55.2

^aSample included 34 *K. pneumoniae*, 1 *K. oxytoca*, 1 *E. coli*, and 2 *P. aeruginosa* isolates.^bPatient with OXA-48-producing *E. coli* who had failed imipenem treatment (MIC not reported).

with KPC-producing *K. pneumoniae* with an imipenem MIC of <2 and a meropenem MIC of 2 had failed treatment with a multidrug regimen that included meropenem; carbapenem treatment was not tried in the other two patients with imipenem MICs of 2. Exact results of disk diffusion testing for CAZ-AVI susceptibility were available for all but five isolates, with zone diameters ranging from 21 to 32 mm for *Enterobacteriaceae* and from 20 to 23 mm for *P. aeruginosa*. (FDA breakpoints are 21 mm and 18 mm, respectively [14].)

Patient characteristics and prior treatment. Patients' demographic and clinical characteristics are shown in Table 3. All but two patients had received antibiotics before CAZ-AVI (median, 3 drugs) and had failed treatment. Among those who received prior antibiotics, the median duration of treatment before CAZ-AVI was started was 13 days. The most commonly prescribed agents were tigecycline ($n = 26$), meropenem ($n = 18$), gentamicin ($n = 16$), fosfomycin ($n = 14$), and colistin ($n = 11$).

CAZ-AVI treatment. Characteristics of CAZ-AVI treatment are presented in Table 3. The minimum length of treatment was 3 days. Twenty-four patients (63.2%) were given the standard dose of CAZ-AVI throughout their treatment (2 g ceftazidime–0.5 g avibactam every 8 h). Fourteen patients with renal impairment received adjusted doses. Twenty-five patients (65.8%) were treated concurrently with at least one other antibacterial to which their organism was nonresistant *in vitro*; the most common agents were tigecycline ($n = 11$), amikacin ($n = 9$), and fosfomycin ($n = 4$).

Outcomes. Treatment outcomes are presented in Fig. 1. Twenty-eight patients (73.7%; 95% confidence interval [CI], 56.9 to 86.6%) experienced clinical and/or documented microbiological cure at the end of treatment. All-cause in-hospital mortality was 39.5% (95% CI, 24.0 to 56.6%). Ten patients died during their hospitalization because of treatment failure, such that infection-related mortality was 26.3% (95% CI, 13.4 to 43.1%). Among these 10 patients, the median time from the start of CAZ-AVI treatment to death was 19 days (interquartile range [IQR], 7 to 31 days). Six of them were still receiving CAZ-AVI within 1 day of their death. Nine of the 13 patients (69.2%) who received CAZ-AVI as monotherapy achieved clinical and/or microbiological cure,

TABLE 2 MICs (in $\mu\text{g/ml}$) of carbapenems in 33 isolates for which susceptibility was reported quantitatively

Organism and antibiotic ^a	No. of isolates with MIC of:			
	<2	2	4 to 8	>8
<i>Enterobacteriaceae</i>				
Imipenem ($n = 29$)	1	2	2	24
Meropenem ($n = 27$)	0	1	2	24
<i>P. aeruginosa</i>				
Imipenem ($n = 2$)	0	0	0	2
Meropenem ($n = 1$)	0	0	0	1

^a n , number of isolates.

TABLE 3 Characteristics of patients with carbapenem-resistant infections treated with compassionate-use CAZ-AVI

Characteristic	Value (n = 38) ^a
Demographic characteristics	
Age in yr, median (IQR) ^b	61 (47–67)
Male sex	25 (65.8)
Location before hospitalization	
Home	33 (86.8)
Transferred from another hospital	5 (13.2)
Comorbidities	
Transplant recipient	5 (13.2)
Diabetes mellitus	8 (21.1)
Immunosuppression ^c	10 (26.3)
Renal disease	7 (18.4)
Cardiovascular disease	11 (28.9)
McCabe score of >1	19 (50.0)
Infection characteristics	
Organism and carbapenemase	
<i>Klebsiella pneumoniae</i>	
KPC	22
OXA-48	12
<i>Klebsiella oxytoca</i> (KPC)	1
<i>Escherichia coli</i> (OXA-48)	1
<i>Pseudomonas aeruginosa</i>	2
Hospital-acquired infection	34 (89.5)
Bacteremia	26 (68.4)
Polymicrobial infection	11 (29.0)
Life-threatening infection (high risk of death within 30 days)	23 (60.5)
Antibiotics before CAZ-AVI	
Received antibiotics before CAZ-AVI for this infection	36 (94.7)
Days of antibiotic treatment before CAZ-AVI, median (IQR)	13 (7–31)
No. of antibiotics before CAZ-AVI, median (IQR)	3 (3–4)
Other treatments before CAZ-AVI	
Surgery to remove the source of infection	16 (42.1)
Removal of foreign body involved in infection	9 (23.7)
Clinical status at start of CAZ-AVI treatment	
Mechanical ventilation	14 (36.8)
Vasopressor support	17 (44.7)
Unconscious	12 (31.6)
CAZ-AVI treatment	
Days of treatment, median (IQR)	16 (14–21)
Extended infusion	36 (94.7)
Concurrent antibiotic treatment ^d	25 (65.8)
Received standard CAZ-AVI dose	24 (63.2)

^aValues are number (%) of patients unless indicated otherwise.^bIQR, interquartile range.^cImmunosuppression was defined as posttransplant, chemotherapy in past 6 weeks, systemic steroids (>20 mg of prednisone) or other immunosuppressive agents in past 2 weeks, absolute neutrophil count of <500/ μ l, or HIV/AIDS.^dDuring CAZ-AVI treatment, patient received another antibiotic to which organism was nonresistant *in vitro*.

compared to 19 out of 25 patients (76.0%) given a concurrent antibiotic with *in vitro* activity against their pathogen ($P = 0.71$).

For three of the four patients with documented microbiological failure, the repeat positive isolates were tested for susceptibility to CAZ-AVI; none had developed CAZ-AVI resistance. Two patients with documented microbiological cure of infections caused by KPC-producing *K. pneumoniae* experienced a relapse, one at 16 days and the other at 30 days after the end of both CAZ-AVI treatment and discharge from the hospital. In the first case, the isolate remained susceptible to CAZ-AVI by disk diffusion testing and the patient experienced clinical and microbiological cure of the relapse infection following dual therapy with CAZ-AVI and gentamicin. In the second case, the relapse isolate was




FIG 1 Outcomes of patients with carbapenem-resistant infections treated with compassionate-use CAZ-AVI.

not tested for CAZ-AVI susceptibility; repeat CAZ-AVI treatment was not considered because of the patient's poor prognosis, and the patient died 73 days after infection onset of causes not directly related to the infection.

Table 4 presents treatment outcomes according to infection site. Among the 38 patients, there were 46 infections, not including secondary bacteremia. Eighteen of the 46 infections (39.1%) were either cIAI or cUTI, indications for which CAZ-AVI was approved; 13 patients had only cIAI or cUTI, with or without bacteremia. All-cause mortality was 14.3% in patients with primary or central-line associated bacteremia and 42.3% in patients with any bacteremia. Five out of 24 patients (20.8%) with documented microbiological cure died (of causes unrelated to the infection), whereas 10 of 14 patients (71.4%) with no documented microbiological cure died ($P = 0.01$).

Table 5 compares characteristics of patients by treatment outcome. Patients treated for a longer time with other antibiotics prior to CAZ-AVI administration were less likely to experience clinical cure ($P = 0.06$) and microbiological cure ($P = 0.01$). Among patients infected with *Enterobacteriaceae*, survival was higher in patients with KPC carbapenemase than in those with OXA-48: 17 of 23 patients (73.9%) with KPC-producing organisms survived until discharge, compared to 5 of 13 patients (38.5%)

TABLE 4 Outcomes of patients with carbapenem-resistant infections treated with compassionate-use CAZ-AVI, by infection site

Infection site ^a	Total no. of cases	No. (%) of cases with:			Patients with:				Mortality among patients with microbiological cure	
		Bacteremia	Life-threatening infection	Documented microbiological cure	Clinical cure		In-hospital death		No. (%)	95% CI
					No. (%)	95% CI	No. (%)	95% CI		
All patients	38	26 (68.4)	23 (60.5)	24 (63.2)	26 (68.4)	51.3–82.5	15 (39.5)	24.0–56.6	5 (20.8)	7.1–42.2
Intra-abdominal	15	11 (73.3)	8 (53.3)	6 (40.0)	10 (66.7)	38.4–88.2	6 (40.0)	16.3–67.7	1 (16.7)	0.4–64.1
Pneumonia ^b	7	6 (85.7)	5 (71.4)	3 (42.9)	3 (42.9)	9.9–81.6	5 (71.4)	29.0–96.3	1 (33.3)	0.8–90.6
Skin and soft tissue	4	3 (75.0)	1 (25.0)	1 (25.0)	1 (25.0)	0.6–80.6	2 (50.0)	6.8–93.2	0 (0.0)	0.0–97.5
Urinary tract	3	2 (66.7)	1 (33.3)	2 (66.7)	2 (66.7)	9.4–99.2	2 (66.7)	9.4–99.2	1 (50)	1.3–98.7
Primary or catheter-associated bacteremia	7	7 (100)	7 (100)	7 (100.0)	7 (100)	59.0–100	1 (14.3)	0.4–57.9	1 (14.3)	0.4–57.9
Any bacteremia	26	26 (100)	20 (76.9)	18 (69.2)	18 (69.2)	48.2–85.7	11 (42.3)	23.4–63.1	4 (22.2)	6.4–47.6
Endocarditis	2	1 (50.0)	1 (50.0)	2 (100.0)	2 (100.0)	15.8–100	1 (50.0)	1.3–98.7	1 (50)	1.3–98.7
Osteomyelitis	3	0 (0.0)	0 (0.0)	2 (66.7)	2 (66.7)	9.4–99.2	1 (33.3)	0.8–90.6	0 (0.0)	0.0–84.2
Surgical site infection	2	1 (50.0)	2 (100)	1 (50.0)	1 (50.0)	1.3–98.7	1 (50.0)	1.3–98.7	0 (0.0)	0.0–97.5
Other ^c	3	1 (33.3)	2 (66.7)	3 (100)	2 (66.7)	9.4–99.2	1 (33.3)	0.8–90.6	1 (33.3)	0.8–90.6

^aPatients may have multiple infection sites.

^bPneumonia cases included 6 cases of ventilator-associated pneumonia and 1 case of hospital-acquired pneumonia.

^cOther infection types (1 patient each) were ventriculitis/subdural abscess, prosthetic joint infection, and mucositis.

TABLE 5 Comparison of characteristics of patients with carbapenem-resistant infections treated with compassionate-use CAZ-AVI, by treatment outcome^a

Characteristic	Clinical cure		Documented microbiological cure		Survival to hospital discharge		P value
	Yes (n = 26)	No (n = 12)	Yes (n = 24)	No (n = 14)	Yes (n = 23)	No (n = 15)	
Age in yr, median (IQR)	54 (46–65)	62 (58–69)	63 (48–70)	58 (46–63)	54 (47–66)	61 (44–68)	0.68
Male sex	17 (65.4)	8 (66.7)	14 (58.3)	11 (78.6)	16 (69.6)	9 (60.0)	0.73
McCabe score of > 1	14 (53.9)	5 (41.7)	14 (58.3)	5 (35.7)	12 (52.2)	7 (46.7)	1.00
Days of antibiotic treatment before CAZ-AVI, median (IQR)	9 (5–17)	21 (10–50)	8 (3–16)	29 (11–45)	10 (5–20)	15 (6–31)	0.38
No. of antibiotics before CAZ-AVI, median (IQR)	3 (2–3)	3 (3–6)	3 (2–3)	3 (3–6)	3 (2–4)	3 (2–4)	0.75
Life-threatening infection	15 (57.7)	8 (66.7)	14 (58.3)	9 (64.3)	12 (52.2)	11 (73.3)	0.31
Carbapenemase ^b							
KPC	17 (68.0)	6 (54.6)	18 (75.0)	5 (41.7)	17 (77.3)	6 (42.9)	0.07
OXA-48	8 (32.0)	5 (45.5)	6 (25.0)	7 (58.3)	5 (22.7)	8 (57.1)	
Polymicrobial infection	6 (23.1)	5 (41.7)	4 (16.7)	7 (50.0)	6 (26.1)	5 (33.3)	0.72
cIAI or CUTI as sole infection site (with or without bacteremia)	10 (38.5)	3 (25.0)	7 (29.2)	6 (42.9)	9 (39.1)	4 (26.7)	0.50
Bacteremia	18 (69.2)	8 (66.7)	18 (75.0)	8 (57.1)	15 (65.2)	11 (73.3)	0.73
Surgery to remove the source of infection before or during CAZ-AVI	13 (50.0)	6 (50.0)	11 (45.8)	8 (57.1)	12 (52.2)	7 (46.7)	1.00
Mechanical ventilation at start of CAZ-AVI	8 (30.8)	6 (50.0)	8 (33.3)	6 (42.9)	5 (21.7)	9 (60.0)	0.04
On vasopressors at start of CAZ-AVI	11 (42.3)	6 (50.0)	10 (41.7)	7 (50.0)	8 (34.8)	9 (60.0)	0.19
Unconscious at start of CAZ-AVI	6 (23.1)	6 (50.0)	6 (25.0)	6 (42.9)	4 (17.4)	8 (53.3)	0.03
Concurrent active antibiotic treatment ^c	18 (69.2)	7 (58.3)	16 (66.7)	9 (64.3)	14 (60.9)	11 (73.3)	0.50
Received standard CAZ-AVI dose	17 (65.4)	7 (58.3)	15 (62.5)	9 (64.3)	15 (65.2)	9 (60.0)	1.00

^aValues are number (%) of patients unless indicated otherwise.^bAmong 36 patients with *Enterobacteriaceae* infections.^cDuring CAZ-AVI treatment, patient received another antibiotic to which the organism was nonresistant *in vitro*.

with OXA-48 producers ($P = 0.07$). These 2 groups of patients did not differ by the proportion with life-threatening infection ($P = 0.73$) or high McCabe score ($P = 0.50$). Neither of the two cases of relapse occurred in patients with OXA-48; one of the four cases of documented microbiological failure occurred in a patient with OXA-48.

Six patients (15.8%) developed adverse events that were attributed to CAZ-AVI. Blood alkaline phosphatase increased in two patients; nausea/vomiting, *Clostridium difficile*-associated diarrhea, convulsions, and disorientation progressing to stupor occurred in one patient each.

DISCUSSION

The proportion of Gram-negative infections caused by carbapenem-resistant strains is increasing. According to U.S. surveillance systems, carbapenem resistance in nosocomial infections caused by *Klebsiella* spp. rose from 1.6% in 2001 to 10.4% in 2011 (15). Regional differences in resistance are striking: in the 2014 report of the EARS-Net European surveillance system, the proportion of *Klebsiella* spp. isolated from blood or cerebrospinal fluid that was carbapenem resistant ranged from 0.0% in Norway, Sweden, Finland, and Estonia to 62.3% in Greece (16).

Treatment options for carbapenem-resistant organisms are limited; they include primarily colistin, aminoglycosides, tigecycline (for CRE only), fosfomycin, and double-carbapenem therapy. A recent systematic review by Falagas et al. of 20 nonrandomized studies compared mortality following different antibiotic regimens for CRE infections (17). Mortality was variously defined as 28- or 30-day mortality, in-hospital mortality, infection-related mortality, or unspecified. Mortality associated with the most common treatment regimens was up to 57% for colistin alone, up to 80% for tigecycline alone, up to 64% for colistin-tigecycline, up to 50% for gentamicin-tigecycline, and up to 67% for combined therapy with colistin and a carbapenem. While colistin has been considered the mainstay of therapy for carbapenem-resistant infections (18), colistin resistance among CRE isolates has increased (19). In our study, 59% of tested isolates were colistin resistant. Although we lack information about the laboratory methods used at each study site, and some methods for colistin susceptibility testing are flawed (with inaccurate results more likely to be false-susceptible than false-resistant) (20), these test results are relevant because they influenced the decision to use CAZ-AVI. The rationale behind double-carbapenem therapy for carbapenem-resistant organisms is that ertapenem, which is more easily hydrolyzed by carbapenemases, saturates these enzymes so that higher concentrations of the second carbapenem are available to treat the infection. Two recent articles summarized the outcomes of 29 patients who received double-carbapenem therapy for infections caused by carbapenem-resistant *K. pneumoniae*; clinical success was achieved in 16 cases (55%) (21, 22).

In our study, 73.7% of patients with infections caused by CRE or CRPa who received salvage therapy with CAZ-AVI experienced clinical and/or microbiological cure, and all-cause in-hospital mortality was 39.5%. Among those cured were patients with infections at difficult-to-treat sites, such as endocarditis and osteomyelitis, for which CAZ-AVI has not been studied. Mortality among patients with bacteremia was 42%. In previous studies, all-cause mortality from CRE bacteremia ranged from 19% (among patients given combination therapy that included a carbapenem) to 94% (23–28). In our study, microbiological cure was a predictor of survival: 79% of patients with negative cultures at the end of treatment survived until discharge. Among the 21% of patients who achieved microbiological cure but did not survive, mortality was attributed to other, non-infection-related causes. The delayed onset of CAZ-AVI treatment was associated with worse clinical and microbiological outcomes. Minimizing CAZ-AVI use in order to prevent the emergence of resistance is critical; however, waiting to exhaust all other (and potentially more toxic) treatment options before resorting to CAZ-AVI may reduce a patient's likelihood of being cured.

We found that having an infection caused by an OXA-48-producing pathogen, as opposed to a KPC-producing pathogen, was a predictor of mortality; the association did not reach statistical significance ($P = 0.07$), likely because of the small sample size.

OXA-48 does not hydrolyze ceftazidime efficiently and does not usually cause ceftazidime resistance. However, most OXA-48-producing isolates are resistant to ceftazidime due to coproduction of ESBL enzymes (29). Because ESBLs are well inhibited by avibactam and OXA-48 is partly inhibited by avibactam (30), the higher mortality observed in patients infected with OXA-48 producers is unexpected and intriguing. We do not have further details on the strains that can shed light on this finding, and further studies are warranted.

Shields et al. recently published a single-center case series of 37 patients with CRE infections treated with CAZ-AVI for at least 3 days (31). In contrast to our study, in which 95% of patients received CAZ-AVI as salvage therapy, CAZ-AVI was the first drug used to treat CRE infections in Shields' study. CAZ-AVI was administered as monotherapy in 70% of patients. Fifty-nine percent of patients experienced clinical success, defined as survival without recurrence at 30 days, clinical improvement, and negative cultures within 7 days of the start of treatment. Thirty-day all-cause mortality was 24%. Because of differences in how outcomes were defined, the results of Shields' study and ours are not directly comparable. Notably, three patients in Shields' study developed CAZ-AVI resistance following 10 to 19 days of treatment. A study of KPC-producing *Enterobacteriaceae* demonstrated the *in vitro* selection of CAZ-AVI-resistant mutants, primarily via alterations to the *bla*_{KPC} Ω loop (32). In our study, when CAZ-AVI susceptibility testing was repeated in 1 of 2 patients with relapse of infection and 3 of 4 patients with documented microbiological failure, no resistance was detected.

We found two additional clinical studies of CAZ-AVI treatment for carbapenem-resistant infections. The first was a case report of a 64-year-old woman who received compassionate-use CAZ-AVI for bacteremia caused by KPC-producing *K. pneumoniae* that did not respond to colistin and dual-carbapenem therapy. She was treated successfully with CAZ-AVI and ertapenem (22). The second was a case series of three patients (aged 72 to 89 years) with CRE bacteremia that had not responded to previous antibiotics. All three achieved clinical and microbiological cure following monotherapy with CAZ-AVI (33).

In our case series, there were six adverse events that were attributed to CAZ-AVI, three of which were severe. One patient who received CAZ-AVI in combination with two other antibiotics developed *C. difficile* infection during treatment. Like nearly all antibiotics, CAZ-AVI alters the normal flora of the colon, predisposing patients to *C. difficile* infection, as has been shown in healthy volunteers given CAZ-AVI (34). The risk of *C. difficile* infection following CAZ-AVI treatment appears to be low: only three of 1,204 (0.2%) patients treated with CAZ-AVI in three phase III clinical trials developed *C. difficile* infection (9–11). There were two neurological adverse events in our case series: convulsions and disorientation with progression to stupor. While it is not clear if these events were related to CAZ-AVI use or to other factors, the neurotoxic effects of cephalosporins are described mostly in the elderly and in patients with renal impairment or prior neurologic disease (35). Both patients in our study who developed neurotoxicity were aged 70 years or over, with normal renal function.

The main limitation of this study, as with any case series, is the lack of a concurrent control group; comparison is to the published experience summarized above. Comparison to a concurrent control group is problematic because of the potential for selection bias, confounding by indication, and confounding by time to treatment. A second limitation is that the experience with CAZ-AVI in the context of compassionate use may not be generalizable to the population of patients who may be candidates for CAZ-AVI treatment. Patients in compassionate-use programs often have severe acute illness or comorbidities that make them ineligible for clinical trials, increasing their risk of adverse outcomes (36). On the other hand, a sample of such patients may be biased toward clinical success: physicians may be more likely to request compassionate-use drugs for patients with a favorable underlying diagnosis, and immortal time bias may be present, as patients need to survive long enough for the drug to arrive. A third limitation is that two-thirds of patients received CAZ-AVI concurrently with other antibiotics with *in vitro* activity against their pathogen. In these patients, it is difficult to

ascertain whether the clinical success was associated with CAZ-AVI, the concurrent antibiotic, or the combination of both. However, when we compared patients who received CAZ-AVI monotherapy to those who were given additional antibiotics, the proportions cured were similar; thus, we believe CAZ-AVI had a major role in the cure. Finally, the small sample size may have precluded us from identifying significant predictors of clinical cure, microbiological cure, and mortality.

In summary, we have presented a case series of patients treated with CAZ-AVI for carbapenem-resistant infections on a compassionate-use basis. The majority of patients had life-threatening infections, some in difficult-to-cure sites, and 95% of them had failed previous antibiotic treatments. Three-quarters of the patients experienced clinical and/or microbiological cure following CAZ-AVI treatment. CAZ-AVI shows promising clinical results for infections for which treatment options are extremely limited.

MATERIALS AND METHODS

Data collection. When data for this paper were collected, CAZ-AVI was not approved for use in the European Union. Upon our request, AstraZeneca (a codeveloper of CAZ-AVI) contacted physicians to whom they had provided compassionate-access CAZ-AVI to treat carbapenem-resistant infections and invited them to participate in the study. AstraZeneca had no further involvement. Twenty-five physicians agreed to participate and were sent case report forms. To be eligible for inclusion, patients must have received at least one dose of CAZ-AVI in the context of compassionate use. Data collected included patient demographic characteristics, comorbidities and McCabe score (37), a description of the infection treated by CAZ-AVI (including an antibiogram), treatment with other antibiotics before or concurrently with CAZ-AVI, reasons for using CAZ-AVI, details of CAZ-AVI treatment, adverse events, the clinical and microbiological response, and relapse of infection.

Microbiological methods. Each hospital conducted antibiotic susceptibility testing according to its own protocols. We report susceptibility as it was interpreted by the local laboratories. All sites tested for susceptibility to CAZ-AVI by disk diffusion; results were interpreted according to the breakpoints set by the FDA (14). Plasma concentrations of ceftazidime and avibactam were not assessed during the course of treatment.

Statistical methods. The main outcome variables were clinical response at the end of treatment, microbiological response at the end of treatment, and all-cause in-hospital mortality. Clinical response was classified as cure, partial improvement, or treatment failure resulting in death and was analyzed as cure versus partial improvement or treatment failure. Microbiological response was classified as a negative culture, positive culture, or culture not repeated and was analyzed as documented cure (negative culture) versus positive culture or culture not repeated. Patient characteristics measured as continuous variables were summarized by median and interquartile range; categorical variables were summarized as proportions. The McCabe score was treated as a dichotomous variable, with a score of >1 indicating underlying disease with death expected within 5 years. Patient characteristics were compared by outcome using the Wilcoxon rank sum test for continuous variables and Fisher's exact test for categorical variables. Analyses were performed using Stata version 13 (Stata Corporation, College Station, TX).

ACKNOWLEDGMENTS

E.T. and Y.C. were responsible for the study design, literature search, data analysis, and writing of the article. All other authors contributed cases to the case series, assisted with data analysis and writing, and reviewed the manuscript.

AstraZeneca provided the list of physicians who contributed cases to this study; AstraZeneca did not initiate, design, or fund this study, nor was it involved in data analysis or writing of the article. J.T.-C. has received unrestricted grants for educational activities from AstraZeneca. N.B. has received honoraria from AstraZeneca for development of educational presentations, consultancy tasks, and/or for the payment of travel/accommodations for scientific purposes. Y.C. has received funds from AstraZeneca in the form of consulting fees, research grants, and speaker's honoraria. All other authors have no interests to declare.

This work was not supported by any external funding agency. Coauthors J.T.-C. and N.B. are funded by the Ministerio de Economía y Competitividad, Instituto de Salud Carlos III, cofinanced by the European Development Regional Fund (ERDF) "A way to achieve Europe," Spanish Network for Research in Infectious Diseases (REIPI RD12/0015).

REFERENCES

1. U.S. Food and Drug Administration. 26 February 2015. FDA approves new antibacterial drug Avycaz. FDA news release. U.S. FDA, Silver Spring, MD. <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm435629.htm>. Accessed 13 September 2016.
2. Zhanel GG, Lawson CD, Adam H, Schweizer F, Zelenitsky S, Lagacé-Wiens PR, Denisuk A, Rubinstein E, Gin AS, Hoban DJ, Lynch JP, III, Karlowsky JA. 2013. Ceftazidime-avibactam: a novel cephalosporin/ β -lactamase inhibitor combination. *Drugs* 73:159–177. <https://doi.org/10.1007/s40265-013-0013-7>.
3. van Duin D, Bonomo RA. 2016. Ceftazidime/avibactam and ceftolozane/tazobactam: second-generation β -lactam/ β -lactamase inhibitor combinations. *Clin Infect Dis* 63:234–241. <https://doi.org/10.1093/cid/ciw243>.
4. Castanheira M, Mills JC, Costello SE, Jones RN, Sader HS. 2015. Ceftazidime-avibactam activity tested against *Enterobacteriaceae* isolates from U.S. hospitals (2011 to 2013) and characterization of β -lactamase-producing strains. *Antimicrob Agents Chemother* 59:3509–3517. <https://doi.org/10.1128/AAC.00163-15>.
5. Sader HS, Castanheira M, Farrell DJ, Flamm RK, Jones RN. 2015. Ceftazidime-avibactam activity when tested against ceftazidime-nonsusceptible *Citrobacter* spp., *Enterobacter* spp., *Serratia marcescens*, and *Pseudomonas aeruginosa* from United States medical centers (2011–2014). *Diagn Microbiol Infect Dis* 83:389–394. <https://doi.org/10.1016/j.diagmicrobio.2015.06.008>.
6. Dupont H, Gaillot O, Goetgheluck AS, Plassart C, Emond JP, Lecuru M, Gaillard N, Derdouri S, Lemaire B, Girard de Courtilles M, Cattoir V, Mammeri H. 2015. Molecular characterization of carbapenem-nonsusceptible enterobacterial isolates collected during a prospective interregional survey in France and susceptibility to the novel ceftazidime-avibactam and aztreonam-avibactam combinations. *Antimicrob Agents Chemother* 60:215–221. <https://doi.org/10.1128/AAC.01559-15>.
7. Vazquez JA, González Patzán LD, Stricklin D, Duttaroy DD, Kreidly Z, Lipka J, Sable C. 2012. Efficacy and safety of ceftazidime-avibactam versus imipenem-cilastatin in the treatment of complicated urinary tract infections, including acute pyelonephritis, in hospitalized adults: results of a prospective, investigator-blinded, randomized study. *Curr Med Res Opin* 28:1921–1931. <https://doi.org/10.1185/03007995.2012.748653>.
8. Lucasti C, Popescu I, Ramesh MK, Lipka J, Sable C. 2013. Comparative study of the efficacy and safety of ceftazidime/avibactam plus metronidazole versus meropenem in the treatment of complicated intra-abdominal infections in hospitalized adults: results of a randomized, double-blind, phase II trial. *J Antimicrob Chemother* 68:1183–1192. <https://doi.org/10.1093/jac/dks523>.
9. Mazuski JE, Gasink LB, Armstrong J, Broadhurst H, Stone GG, Rank D, Llorens L, Newell P, Pachi J. 2016. Efficacy and safety of ceftazidime-avibactam plus metronidazole versus meropenem in the treatment of complicated intra-abdominal infection—results from a randomized, controlled, double-blind, phase 3 program. *Clin Infect Dis* 62:1380–1389. <https://doi.org/10.1093/cid/ciw133>.
10. Wagenlehner FM, Sobel JD, Newell P, Armstrong J, Huang X, Stone GG, Yates K, Gasink LB. 2016. Ceftazidime-avibactam versus doripenem for the treatment of complicated urinary tract infections, including acute pyelonephritis: RECAPTURE, a phase 3 randomized trial program. *Clin Infect Dis* 63:754–762. <https://doi.org/10.1093/cid/ciw378>.
11. Carmeli Y, Armstrong J, Laud PJ, Newell P, Stone G, Wardman A, Gasink LB. 2016. Ceftazidime-avibactam or best available therapy in patients with ceftazidime-resistant *Enterobacteriaceae* and *Pseudomonas aeruginosa* complicated urinary tract infections or complicated intra-abdominal infections (REPRISE): a randomised, pathogen-directed, phase 3 study. *Lancet Infect Dis* 16:661–673. [https://doi.org/10.1016/S1473-3099\(16\)30004-4](https://doi.org/10.1016/S1473-3099(16)30004-4).
12. Mendes RE, Castanheira M, Woosley LN, Costello SE, Stone GG, Flamm RK, Jones RN. 2015. β -Lactamase characterization of baseline *Enterobacteriaceae* (ENT) from a phase 3 trial of ceftazidime-avibactam (CAZ-AVI) for the treatment of infections caused by CAZ-nonsusceptible (NS) pathogens, abstr 1176. Abstr IDWeek 2015, San Diego, CA. <https://idsa.confex.com/idsa/2015/webprogram/Paper53048.html>. Accessed 13 September 2016.
13. Gonzalez-Padilla M, Torre-Cisneros J, Rivera-Espinar F, Pontes-Moreno A, López-Cerero L, Pascual A, Natera C, Rodríguez M, Salcedo I, Rodríguez-López F, Rivero A, Rodríguez-Baño J. 2015. Gentamicin therapy for sepsis due to carbapenem-resistant and colistin-resistant *Klebsiella pneumoniae*. *J Antimicrob Chemother* 70:905–913. <https://doi.org/10.1093/jac/dku432>.
14. Actavis. 2015. Avycaz package insert. Actavis, Parsippany, NJ. http://www.allergan.com/assets/pdf/avycaz_pi. Accessed 13 September 2016.
15. Centers for Disease Control and Prevention. 2013. Vital signs: carbapenem-resistant *Enterobacteriaceae*. *MMWR Morb Mortal Wkly Rep* 62:165–170.
16. European Centre for Disease Prevention and Control. 2016. Antimicrobial resistance interactive database (EARS-Net). http://ecdc.europa.eu/en/healthtopics/antimicrobial-resistance-and-consumption/antimicrobial_resistance/database/Pages/database.aspx. Accessed 13 September 2016.
17. Falagas ME, Lourida P, Poulikakos P, Rafailidis PI, Tansarli GS. 2014. Antibiotic treatment of infections due to carbapenem-resistant *Enterobacteriaceae*: systematic evaluation of the available evidence. *Antimicrob Agents Chemother* 58:654–663. <https://doi.org/10.1128/AAC.01222-13>.
18. Pogue JM, Ortwin JK, Kaye KS. 2015. Optimal usage of colistin: are we any closer? *Clin Infect Dis* 61:1778–1780. <https://doi.org/10.1093/cid/civ723>.
19. van Duin D, Doi Y. 2015. Outbreak of colistin-resistant, carbapenemase-producing *Klebsiella pneumoniae*: are we at the end of the road? *J Clin Microbiol* 53:3116–3117. <https://doi.org/10.1128/JCM.01399-15>.
20. Dafopoulou K, Zarkotou O, Dimitroulia E, Hadjichristodoulou C, Gennimata V, Pournaras S, Tsakris A. 2015. Comparative evaluation of colistin susceptibility testing methods among carbapenem-nonsusceptible *Klebsiella pneumoniae* and *Acinetobacter baumannii* clinical isolates. *Antimicrob Agents Chemother* 59:4625–4630. <https://doi.org/10.1128/AAC.00868-15>.
21. Cprek JB, Gallagher JC. 2015. Ertapenem-containing double-carbapenem therapy for treatment of infections caused by carbapenem-resistant *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 60:669–673. <https://doi.org/10.1128/AAC.01569-15>.
22. Camargo JF, Simkins J, Beduschi T, Tekin A, Aragon L, Pérez-Cardona A, Prado CE, Morris MI, Abbo LM, Cantón R. 2015. Successful treatment of carbapenemase-producing pandrug-resistant *Klebsiella pneumoniae* bacteremia. *Antimicrob Agents Chemother* 59:5903–5908. <https://doi.org/10.1128/AAC.00655-15>.
23. Villegas MV, Pallares CJ, Escandón-Vargas K, Hernández-Gómez C, Correa A, Álvarez C, Rosso F, Matta L, Luna C, Zurita J, Mejía-Villatoro C, Rodríguez-Noriega E, Seas C, Cortesía M, Guzmán-Suárez A, Guzmán-Blanco M. 2016. Characterization and clinical impact of bloodstream infection caused by carbapenemase-producing *Enterobacteriaceae* in seven Latin American countries. *PLoS One* 11:e0154092. <https://doi.org/10.1371/journal.pone.0154092>.
24. Balkan I, Aygün G, Aydın S, Mutcalı SI, Kara Z, Kuşkuç M, Midilli K, Şemen V, Aras S, Yemişen M, Mete B, Özars R, Saltoğlu N, Tabak F, Öztürk R. 2014. Blood stream infections due to OXA-48-like carbapenemase-producing *Enterobacteriaceae*: treatment and survival. *Int J Infect Dis* 26:51–56. <https://doi.org/10.1016/j.ijid.2014.05.012>.
25. Ben-David D, Kordevani R, Keller N, Tal I, Marzel A, Gal-Mor O, Maor Y, Rahav G. 2012. Outcome of carbapenem resistant *Klebsiella pneumoniae* bloodstream infections. *Clin Microbiol Infect* 18:54–60. <https://doi.org/10.1111/j.1469-0691.2011.03478.x>.
26. Chang HJ, Hsu PC, Yang CC, Kuo AJ, Chia JH, Wu TL, Lee MH. 2011. Risk factors and outcomes of carbapenem-nonsusceptible *Escherichia coli* bacteremia: a matched case-control study. *J Microbiol Immunol Infect* 44:125–130. <https://doi.org/10.1016/j.jmii.2010.06.001>.
27. Tumbarello M, Trecarichi EM, De Rosa FG, Giannella M, Giacobbe DR, Bassetti M, Losito AR, Bartoletti M, Del Bono V, Corcione S, Maiuro G, Tedeschi S, Celani L, Cardellino CS, Spanu T, Marchese A, Ambretti S, Cauda R, Viscoli C, Viale P, ISGRI-SITA (Italian Study Group on Resistant Infections of the Società Italiana Terapia Antinfettiva). 2015. Infections caused by KPC-producing *Klebsiella pneumoniae*: differences in therapy and mortality in a multicentre study. *J Antimicrob Chemother* 70:2133–2143. <https://doi.org/10.1093/jac/dkv086>.
28. Daikos GL, Tsaousi S, Tzouveleakis LS, Anyfantis I, Psychogiou M, Argyropoulou A, Stefanou I, Sympa V, Miriagou V, Nepka M, Georgiadou S, Markogiannakis A, Goukos D, Skoutelis A. 2014. Carbapenemase-producing *Klebsiella pneumoniae* bloodstream infections: lowering mortality by antibiotic combination schemes and the role of carbapenems. *Antimicrob Agents Chemother* 58:2322–2328. <https://doi.org/10.1128/AAC.02166-13>.

29. Poirel L, Potron A, Nordmann P. 2012. OXA-48-like carbapenemases: the phantom menace. *J Antimicrob Chemother* 67:1597–1606. <https://doi.org/10.1093/jac/dks121>.
30. Lagacé-Wiens P, Walkty A, Karlowsky JA. 2014. Ceftazidime-avibactam: an evidence-based review of its pharmacology and potential use in the treatment of Gram-negative bacterial infections. *Core Evid* 9:13–25. <https://doi.org/10.2147/CE.S40698>.
31. Shields RK, Potoski BA, Haidar G, Hao B, Doi Y, Chen L, Press EG, Kreiswirth BN, Clancy CJ, Nguyen MH. 13 September 2016. Clinical outcomes, drug toxicity and emergence of ceftazidime-avibactam resistance among patients treated for carbapenem-resistant Enterobacteriaceae infections. *Clin Infect Dis pii: ciw636*. Epub ahead of print.
32. Livermore DM, Warner M, Jamroz D, Mushtaq S, Nichols WW, Mustafa N, Woodford N. 2015. In vitro selection of ceftazidime-avibactam resistance in *Enterobacteriaceae* with KPC-3 carbapenemase. *Antimicrob Agents Chemother* 59:5324–5330. <https://doi.org/10.1128/AAC.00678-15>.
33. Wu G, Abraham T, Lee S. 2016. S. Ceftazidime-avibactam for treatment of carbapenem-resistant *Enterobacteriaceae* bacteremia. *Clin Infect Dis* 63: 1147–1148. <https://doi.org/10.1093/cid/ciw491>.
34. Rashid MU, Rosenborg S, Panagiotidis G, Löfdal KS, Weintraub A, Nord CE. 2015. Ecological effect of ceftazidime/avibactam on the normal human intestinal microbiota. *Int J Antimicrob Agents* 46:60–65. <https://doi.org/10.1016/j.ijantimicag.2015.02.027>.
35. Grill MF, Maganti R. 2008. Cephalosporin-induced neurotoxicity: clinical manifestations, potential pathogenic mechanisms, and the role of electroencephalographic monitoring. *Ann Pharmacother* 42:1843–1850. <https://doi.org/10.1345/aph.1L307>.
36. Rosenblatt M, Kuhlik B. 2015. Principles and challenges in access to experimental medicines. *JAMA* 313:2023–2024. <https://doi.org/10.1001/jama.2015.4135>.
37. McCabe WR, Jackson GG. 1962. Gram negative bacteremia. I. Etiology and ecology. *Arch Intern Med* 110:845–847.

From: pfizer@copyright.com
To: [Fandino, Cecil](#)
Subject: [EXTERNAL] Requested copyright permission granted on order 12139299 - Antimicrobial agents and chemotherapy, 2017
Date: martes, 8 de octubre de 2019 19:32:42



Dear Cecil,

Your copyright permission request has been approved; you may now proceed with your selected usage scenario for this order: Ordering 1 copy for the following stated purpose: "EXTERNAL: Reactively distribute a single digital copy to an external customer or Health Care Professional (HCP), prospect or client in response to an informational inquiry".

If you need to purchase additional permissions; please visit "My Request History," select the order, and click the "Re-Order" button.

Citation Information:

Title:
Publication Name: Antimicrobial agents and chemotherapy
ISSN/ ISBN: 00664804
Year: 2017
Vol (Iss) Pgs: () p.
Supplement:
Author(s):

Additional Info:

Order Info Link:
<https://www.rightfind.com/vlib/order/orderinfo.aspx?ClientId=65&Id=43d06091-1332-45ac-bbd6-89308c8ed46c&lang=en-us>
Cost Center: 0003035023
Company: PFIZER
PO:
PI Order number(optional):

Order Details:

Order ID: 12139299
Client ID: 65
Urgency: Normal
Type: CR Perm (Journal Article)
Number of Copies: 1
Ordered For:
Cecil.Fandino@pfizer.com
Usage: Ordering 1 copy for the following stated purpose: "EXTERNAL: Reactively distribute a single digital copy to an external customer or Health Care Professional (HCP), prospect or client in response to an informational inquiry"

The contents of the attached document are copyrighted works. You have secured permission to use this document for the following purpose: Ordering 1 copy for the following stated purpose: "EXTERNAL: Reactively distribute a single digital copy to an external customer or Health Care Professional (HCP), prospect or client in response to an informational inquiry". You have not secured permission through Copyright Clearance Center for any other purpose but may have other rights pursuant to other arrangements you may have with the copyright owner or an authorized licensing body. To the extent that a publisher or other appropriate rights-holder has placed additional terms and conditions on your use of this document, such terms and conditions are specified herein under "Copyright Terms". If you need to secure additional permission with respect to this content, please purchase the appropriate permission via RightFind™.

Copyright Terms: This use is only to be utilized by Medical Information (MI) / Medical Professionals who are permitted to provide information (both on-label and off-label) in response to UNSOLICITED MI Inquiries (definition provided below). Any off-label inquiries received by non-medical professionals must be forwarded to these groups for an appropriate response. This right excludes members of Marketing and Sales. Please note that only a single copy of material may be sent per inquiry and that no more than 10 copies of a specific article are supplied for such purposes under this provision. For additional copies, Licensee should ask ASM for reprint rights, which are not covered by this License.

Questions?

Please contact pfizer@copyright.com.

Please include your Order ID in all correspondence.

Copyright 2015 Copyright Clearance Center

This message (including attachments) is confidential, unless marked otherwise. It is intended for the addressee(s) only. If you are not an intended recipient, please delete it without further distribution and reply to the sender that you have received the message in error.